

CRYPTOSPORIDIUM PARVUM AND CRYPTOSPORIDIUM HOMINIS

THE ORGANISM/TOXIN

Cryptosporidium parvum and *Cryptosporidium hominis* are intracellular protozoan parasites that may produce gastrointestinal symptoms when ingested by humans. Up until 2002 *C. parvum* was named *C. parvum* genotype 2 (cattle genotype) and *C. hominis* was named *C. parvum* genotype 1 (the human genotype) (Morgan-Ryan *et al.*, 2002). They are now recognised as different species based on genetic distinctions, but care should be taken when reviewing pre-2002 literature for this reason.

There are a further thirteen recognised species of the genus *Cryptosporidium*. *C. parvum* and *C. hominis* are the most widespread and most frequently associated with human infections.

C. hominis and *C. parvum* are transmitted person-to-person. *C. parvum* is also transmitted animal-to-animal or animal-to-human (zoonotically) (Awad-El Kariem 1999). Both organisms move between hosts in an environmentally-resistant oocyst form that is excreted with faeces. An oocyst is the resting stage of the protozoan, similar to a bacterial spore, and may facilitate transmission of the organism via contaminated water or food.

GROWTH AND CONTROL

N.B. Viability refers to whether the organism is still alive, infectivity relates to the organism's ability to infect a host.

Growth

The organism does not replicate outside the intestines of hosts.

Survival

Temperature

Oocysts remain infective in water at:

- 67.5°C for 1 minute
- -5°C for up to 8 weeks
- -10°C for up to 7 days
- -15°C for up to 24 hours
- -20°C for up to 5 hours.

(Fayer 1994; Fayer and Nerad 1996; Fayer *et al.*, 1998)

Inactivation

Temperature

Oocysts become non-infective when:

- Pasteurised (71.7°C, 15 seconds) (Harp *et al.*, 1996)
- Held in water at 64.2°C for 2 minutes (Fayer 1994)
- Frozen in de-ionised water (-20°C for 24 hours and -70°C for 1 hour) (Fayer and Nerad, 1996)
- Frozen in liquid nitrogen (-196°C) (Robertson *et al.*, 1992)
- Frozen in ice-cream (-20°C for 24 hours) (Deng and Cliver, 1999).

pH

- pH <4 or >11 results in minor viability loss (Jenkins *et al.*, 1998; Kniel *et al.*, 2003).
- Ammonia can enhance inactivation at low and high pH values (Jenkins *et al.*, 1998).
- Fruit juice organic acids inhibit oocyst infectivity (Kniel *et al.*, 2003).

pH

Viable *C. parvum* oocysts were detected after 14 days, in media based on citric acid (pH 4.6, 3.6 and 2.6), acetic acid (pH 3.6) or lactic acid (pH 4.6) held at 4°C or 22°C (Dawson *et al.*, 2004).

Water Activity

Oocysts survive on wet stainless steel (93.1% viable after 30 minutes) (Deng and Cliver, 1999).

Water activity

Oocysts are sensitive to air drying (e.g. at ambient temperatures). Viability on stainless steel after 10 minutes = 81%, 1 hour = 69%, 2 hours = 45%, 4 hours = 5% (Deng and Cliver, 1999).

At a_w 0.85, 99.9% are non-infective after 24 hours at 28°C, or 1 week at 7°C. At a_w 0.95, 99.99% are non-infective after 1 week at 28°C, or 2 weeks at 7°C (Rose and Slifko, 1999).

Radiation

Sensitive to UV from sunlight and UV lamps. Under strong sunlight for 8 and 12 hours, oocyst viability reduced from 98% to 11.7% and 0.3%, respectively. A synergistic effect occurs at temperatures greater than 45°C (Méndez-Hermida *et al.*, 2007).

CRYPTOSPORIDIUM PARVUM AND CRYPTOSPORIDIUM HOMINIS

Disinfectants / Sanitisers

Oocysts are resistant to chlorine and monochloramine (Carey *et al.*, 2004). Swimming pool transmission has been problematic for this reason.

Only 5/35 disinfectants tested were effective with short exposure (O'Donoghue, 1995). These were:

- Ammonia 5% for 120 minutes or 50% for 30 minutes
- Formalin 10% for 120 minutes

- Hydrogen peroxide 10 vol; 3% for 30 minutes
- "Exspor" working dilution 30 minutes
- Oo-cide (ammonia) 5% for 5 minutes.

Ozone (25°C, 1ppm) renders oocysts non-infective after 3 minutes (90% decrease), 5 minutes (99%) and 10 minutes (99.9%) (Korich *et al.*, 1990).

CLINICAL PICTURE

Incubation: Time from ingestion to illness usually 3-5 days, up to 2 weeks (Fayer, 2004).

Symptoms: Copious watery non-bloody diarrhoea, vomiting, anorexia, fever, malaise, abdominal cramping and weight loss. Usually self-limiting. Symptoms typically last 2-4 days but can last up to two weeks. Infectious oocysts usually excreted for 6-9 days, but can be excreted by asymptomatic carriers for up to 2 months or longer following infection. The respiratory system may also be adversely affected by the illness.

Condition: Cryptosporidiosis.

Dose: Healthy human volunteer studies have shown that the infectious dose varies depending on the isolate used. The dose needed to infect 50% of the population ranged from 9-1042 oocysts (Chappell *et al.*, 2006; Okhuysen *et al.*, 1999).

At Risk Groups: Children aged under 1 year and immunocompromised people are more susceptible (Carey *et al.*, 2004).

Long Term Effects: Immunocompromised patients (e.g. AIDS patients or those receiving immunosuppressant drugs) develop severe symptoms that can last months or years and the condition can be fatal. The infection may also spread to other parts of the body in immunocompromised patients, such as the respiratory tract (Ortega, 2007).

Treatment: There is no specific treatment for cryptosporidiosis currently available, but some antibiotics (e.g. spiramycin) are effective (Ortega, 2007).

SOURCES

Human: Person-to-person transmission occurs with *C. parvum* and *C. hominis*. More than 10^8 - 10^{10} oocysts can be excreted daily in the faeces of infected hosts (Fayer *et al.*, 2004).

Animal: *C. parvum* has been identified in 155 species of mammal including domestic animals. Preweaned ruminants, especially calves, are especially vulnerable to infection. Shedding of oocysts increases in calves to a peak at day 12 (excreting approximately 3.89×10^{10} oocysts) (Nydam *et al.*, 2001; Fayer 2004). Flies are potential vehicles of transmission as oocysts can pass unaltered through their gastro-intestinal tract and be deposited in fly faeces (Graczyk *et al.*, 2004; Graczyk *et al.*, 1999), but flies have never been linked to an outbreak. Around 80% of oocysts remain in an infected host's gut and can autoinfect the host. The remaining 20% develop thick (trilaminar) cell walls and are released in the faeces of the infected host into the environment. This can lead to subsequent infection of a new host (O'Donoghue, 1995).

Food: Oocysts have been detected in raw milk, raw meat, raw fruits and vegetables. Foods suspected as acting as vehicles of disease include raw milk, green onions, apple cider, sausage, fruit/vegetables and frozen tripe (Duffy and Moriarty, 2003).

Environment: Oocysts in the environment are particularly resilient, especially under cool, moist conditions. Infectivity can be retained for months, especially in low water temperatures (<5°C but above freezing) (Fayer, 2004). The long term viability of oocysts in animal faeces is unclear. A study of a variety of slurries, soil types, pH ranges and temperatures found that acid to neutral soils at 4°C were most conducive to oocyst survival (still viable after 162 days) (Warnes and Keevil, 2003).

Transmission Routes: Three main routes of transmission are person-to-person, zoonotic, and via faecally-contaminated water or food. Outbreak data for New Zealand suggests that foodborne transmission makes a very small contribution to the overall burden of cryptosporidiosis (Gilbert *et al.*, 2007).

CRYPTOSPORIDIUM PARVUM AND CRYPTOSPORIDIUM HOMINIS

OUTBREAKS AND INCIDENTS

NZ Incidence: 854 cases of cryptosporidiosis were notified in 2009 (19.8 per 100,000 population) and 34 cases were reported as being hospitalised (ESR, 2010b). The highest rates were reported in the West Coast DHB (55.2/100,000; 18 cases) and South Canterbury DHB (50.4/100,000; 28 cases). Of 851 cases, 54.9% were children aged less than 15 years.

New Zealand epidemiological studies

Analysis of 1997-2006 data showed incidence was 2.84 times higher in rural areas (50.68/100,000) than urban areas (17.22/100,000). 55% of cases occurred in spring (Sept-Nov) and 11% occurred summer (Dec-Feb), and the spring peak was most pronounced in rural areas. A small late Summer/early Autumn peak was observed in urban areas in 1998, 1999 and 2001 (Snel *et al.*, 2009).

New Zealand outbreaks

In 2009 there were 20 outbreaks of cryptosporidiosis, involving 68 cases (ESR, 2010a). No cases were hospitalised. Person-to-person transmission was implicated in 15 of these outbreaks (55 cases).

Overseas outbreaks overseas associated with food/water:

Chicken salad: USA, 15 ill, 2 confirmed (26 exposed). Food handler related (CDC, 1996).

Unpasteurised apple cider: USA, 20 confirmed cases, 11 suspected. Apples washed in well-water (Peng *et al.*, 1997).

Unpasteurised apple cider: USA, 154 cases. Apples collected from ground grazed by livestock (Millard *et al.*, 1994).

Green onions: USA, 54 cases (62 exposed). Food handler related and onions unwashed (CDC, 1998).

Fruits/vegetables: USA, 148 cases. Food handler related. (Quiroz *et al.*, 2000).

Salad vegetables: Denmark, 2005, 99 cases (13 confirmed with *C. hominis* infection). Likely contaminated foods were peeled whole carrots served in water, grated carrots, red peppers. Food handler implicated (Ethelberg *et al.*, 2009).

Salad vegetables: Finland, 2008, 72 cases (4 confirmed, 1 with *C. parvum* infection). Salad implicated by case-control study but cause of contamination not identified (Pönkä *et al.*, 2009).

Drinking water: Milwaukee, USA, 403,000 cases (attack rate 55%). One of largest recorded waterborne outbreaks in USA. Hospitalisation rate estimated at 15%, case fatality rate 0.5%.

Drinking water: Ireland, > 476 cases. Three unrelated drinking water outbreaks. Related to septic tank and blocked drain water entering the drinking water system (Xiao 2002).

Drinking water: Australia, 1998. Two episodes of contaminated water supply in Sydney area. No measurable increase in disease (Anonymous, 1998).

Drinking water: Wales, 2005, 218 cases (*C. hominis*). Absence of effective treatment to remove *Cryptosporidium* oocysts at drinking water treatment plant (Mason *et al.*, 2010).

REFERENCES

- Anonymous (1998) The Sydney water incident: July – September 1998. NSW Public Health Bulletin; 9:91-94.
- Awad-EI Kariem F (1999) Does *Cryptosporidium parvum* have a clonal population structure. Parasitology Today; 15:502-504.
- Carey CM, Lee H, Trevors JT (2004) Biology, persistence and detection of *Cryptosporidium parvum* and *Cryptosporidium hominis* oocyst. Water Research 38:818-862.
- CDC, Center for Disease Control and Prevention (1996) Foodborne outbreak of diarrheal illness associated with *Cryptosporidium parvum* Minnesota 1995. Morbidity and Mortality Weekly Report; 45:783-784.
- CDC, Center for Disease Control and Prevention (1998) Foodborne outbreak of cryptosporidiosis Spokane Washington, 1997. Morbidity and Mortality Weekly Report; 47:565-567.
- Dawson DJ, Samuel CM, Scranage V, Atherton CJ (2004) Survival of *Cryptosporidium* species in environments relevant to foods and beverages. Journal of Applied Microbiology 96:1222-1229.
- Deng MQ, Cliver DO (1999) *Cryptosporidium parvum* studies with dairy products. International Journal of Food Microbiology; 46:113-121.
- Chappell CL, Okhuysen PC, Langer-Curry R, Widmer G, Akiyoshi DE, Tanriverdi S, Tzipori S (2006) *Cryptosporidium hominis*: experimental challenge of healthy adults. American Journal of Tropical Medicine and Hygiene; 75:851-857.
- Duffy G, Moriarty EM (2003) *Cryptosporidium* and its potential as a foodborne pathogen. Animal Health Research Reviews; 4: 95-107.
- ESR (2010a) Annual summary of outbreaks in New Zealand 2009. Report FW10048 for the Ministry of Health. Institute of Environmental Science and Research (ESR) Limited, Porirua. Available at: http://www.surv.esr.cri.nz/surveillance/annual_outbreak.php. Accessed 28 July 2010.

CRYPTOSPORIDIUM PARVUM AND CRYPTOSPORIDIUM HOMINIS

- ESR (2010b) Notifiable and other diseases in New Zealand: 2009 Annual Surveillance Report. Report FW10043 for the Ministry of Health. Institute of Environmental Science and Research (ESR) Limited, Porirua.
Available at: http://www.surv.esr.cri.nz/surveillance/annual_surveillance.php. Accessed 28 July 2010.
- Ethelberg S, Lisby M, Vestergaard LS, Enemark HL, Olsen KE, Stensvold CR, Nielsen HV, Porsbo LJ, Plesner AM, Mølbak K (2009) A foodborne outbreak of *Cryptosporidium hominis* infection. *Epidemiology and Infection*; 137: 348-356.
- Fayer R (1994) Effect of high temperature on infectivity of *Cryptosporidium parvum* oocysts in water. *Applied and Environmental Microbiology*; 60:2732-2735.
- Fayer R (2004) *Cryptosporidium*: a water-borne zoonotic parasite. *Veterinary Parasitology*; 126:37-56.
- Fayer R, Nerad T (1996) Effects of low temperatures on viability of *Cryptosporidium parvum* oocysts. *Applied and Environmental Microbiology*; 62: 431-1433.
- Fayer R, Trout JM, Jenkins MC (1998) Infectivity of *Cryptosporidium parvum* oocysts stored in water at environmental temperatures; 84:1165-1169.
- Fayer R, Dubey JP, Lindsay DS (2004) Zoonotic protozoa: from land to sea. *Trends in Parasitology*; 20:531-536.
- Gilbert S, Lake R, Hudson A, Cressey P (2007) Risk Profile: *Cryptosporidium* spp. in shellfish. Report FW0720 for the New Zealand Food Safety Authority. Institute of Environmental Science and Research (ESR) Limited. Available at: http://www.nzfsa.govt.nz/science/risk-profiles/FW0720_Crypto_in_seafood_Final_April_2008.pdf. Accessed 28 July 2010.
- Graczyk TK, Cranfield MR, Fayer R, Bixler H (1999) House flies (*Musca domestica*) as transport hosts of *Cryptosporidium parvum*. *American Journal of Tropical Medicine and Hygiene*; 61:500-504.
- Graczyk TK, Grimes BH, Knight R, Szostakowska B, Kruminis-Lozowska W, Racewicz M, Tamang L, Dasilva AJ, Myjak P (2004) Mechanical transmission of *Cryptosporidium parvum* oocysts by flies. *Wiadomości parazytologiczne*; 50:243-247. [Article in Polish]
- Harp JA, Fayer R, Pesch BA, Jackson GJ (1996) Effect of pasteurization on infectivity of *Cryptosporidium parvum* oocysts in water and milk. *Applied and Environmental Microbiology*; 62:2866-2868.
- Jenkins MB, Bowman DD, Ghiorse WC (1998) Inactivation of *Cryptosporidium parvum* oocysts by ammonia. *Applied and Environmental Microbiology*; 64:784-788.
- Kniel KE, Sumner SS, Lindsay DA, Hackney CR, Pierson MD, Zajac AM, Golden DA, Fayer R (2003) Effect of organic acids and hydrogen peroxide on *Cryptosporidium parvum* viability in fruit juices. *Journal of Food Protection*; 66:1650-1657.
- Korich DG, Mead JR, Madore MS, Sinclair NA, Sterling CR (1990) Effects of ozone, chlorine dioxide, chlorine and monochloramine on *Cryptosporidium parvum* oocyst viability. *Applied and Environmental Microbiology*; 56:1423-1428.
- Mason BW, Chalmers RM, Carnicer-Pont D, Casemore DP (2010) A *Cryptosporidium hominis* outbreak in north-west Wales associated with low oocyst counts in treated drinking water. *Journal of Water Health*; 8: 299-310.
- Méndez-Hermida F, Ares-Mazás E, McGuigan KG, Boyle M, Sichel C, Fernández-Ibáñez P (2007) Disinfection of drinking water contaminated with *Cryptosporidium parvum* oocysts under natural sunlight and using the photocatalyst TiO₂. *Journal of Photochemistry and Photobiology*; 88:105 – 111.
- Millard PS, Gensheimer KF, Addiss DG, Sosin DM, Beckett GA, Houck-Jankoski A, Hudson A (1994) An outbreak of cryptosporidiosis from fresh-pressed apple cider. *Journal of the American Medical Association*; 272:592-596.
- Morgan-Ryan UM, Fall A, Ward LA, Hijawi N, Sulaiman I, Fayer R, Thompson RC, Olson M, Lal A, Eiao L (2002) *Cryptosporidium hominis* n. sp. (Apicomplexa: Cryptosporidiidae) from *Homo sapiens*. *Journal of Eukaryotic Microbiology*; 49:433-440.
- Nydam DV, Wade SE, Schaaf SL, Mohammed HO (2001) Number of *Cryptosporidium parvum* oocysts or *Giardia* spp cysts shed by dairy calves after natural infection. *American Journal of Veterinary Research*; 62:1612-1615.
- O'Donoghue PJ (1995) *Cryptosporidium* and cryptosporidiosis in man and animals. *International Journal for Parasitology*; 25:139-195.
- Okhuysen PC, Chappell CL, Crabb JH, Sterling CR, DuPont HL (1999) Virulence of three distinct *Cryptosporidium parvum* isolates for healthy adults. *Journal of Infectious Disease*; 180:1275-1281.
- Ortega YR (2007) Protozoan parasites. In: Doyle MP, Beuchat LR (eds) *Food Microbiology: Fundamentals and Frontiers*. ASM Press, Washington DC; p663-681.
- Peng MM, Xiao L, Freeman AR, Arrowood MJ, Escalante AA, Weltman AC, Ong CS, MacKenzie WR, Lal AA, Beard CB (1997) Genetic polymorphism among *Cryptosporidium parvum* isolates: evidence of two distinct human transmission cycles. *Emerging Infectious Diseases*; 3:567-573.
- Pönkä A, Kotilainen H, Rimhanen-Finne R, Hokkanen P, Hänninen ML, Kaarna A, Meri T, Kuusi M (2009) A foodborne outbreak due to *Cryptosporidium parvum* in Helsinki, November 2008. *Eurosurveillance*; 14(issue 28).
- Quiroz ES, Bern C, MacArthur JR, Xiao L, Fletcher M, Arrowood MJ, Shay DK, Levy ME, Glass RI, Lal A (2000) An outbreak of cryptosporidiosis linked to a foodhandler. *Journal of Infectious Diseases*; 181:695-700.
- Robertson LJ, Campbell AT, Smith HV (1992) Survival of *Cryptosporidium parvum* oocysts under various environmental pressures. *Applied and Environmental Microbiology*; 58:3494-3500.
- Rose JB and Slifko TR (1999) *Giardia*, *Cryptosporidium*, and *Cyclospora* and their impact on foods: A review. *Journal of Food Protection*; 62:1059-1070.
- Snel SJ, Baker MG, Venugopal K (2009) The epidemiology of cryptosporidiosis in New Zealand, 1997-2006. *New Zealand Medical Journal*; 122:47-61.
- Warnes SW, Keevil CW (2003) Survival of *Cryptosporidium parvum* in faecal wastes and salad crops, In: Duffy G (ed). *Conference proceedings, Cryptosporidium parvum in Food and Water*, Malahide, Dublin. January 29th 2003. Dublin: Teagasc, National Food Centre: p15.
- Xiao L (2002) Three drinking-water-associated cryptosporidiosis outbreaks, Northern Ireland. *Emerging Infectious Diseases*; 8:631-633.